

PUBLIKATIONEN

Stand: August 2012

1. Ultrasound-guided supraclavicular cannulation of the brachiocephalic vein in infants: a retrospective analysis of a case series.
Breschan C, Platzer M, Jost R, Stettner H, Feigl G, Likar R., Paediatr Anaesth. 2012 Aug 3.
2. Project to reduce preventable hospital transfer from homes for the aged and nursing home. Is it really always necessary?
Kada O, Janig H, Wiedermann W, Hollerer A, Pinter G, Likar R., Pflege Z. 2012 Mar;65(3):176-8.
3. From the nursing home to hospital and back again... A mixed methods study on hospital transfers from nursing homes].
Kada O, Brunner E, Likar R, Pinter G, Leutgeb I, Francisci N, Pfeiffer B, Janig H. Z Evid Fortbild Qual Gesundhwes. 2011;105(10):714-22.
4. Quality of postoperative pain therapy in Austria: national survey of all departments of anesthesiology.
Kinstner C, Likar R, Sandner-Kiesling A, Hutschala D, Pipam W, Gustorff B. Anaesthesia. 2011 Sep;60(9):827-34. doi: 10.1007/s00101-011-1911-2.
5. Clonidine for remifentanil-induced hyperalgesia: a double-blind randomized, placebo-controlled study of clonidine under intra-operative use of remifentanil in elective surgery of the shoulder.
Schlimp CJ, Pipam W, Wolrab C, Ohner C, Kager HI, Likar R., Schmerz. 2011 Jun;25(3):290-5.
6. Consecutive, prospective case series of a new method for ultrasound-guided supraclavicular approach to the brachiocephalic vein in children.
Breschan C, Platzer M, Jost R, Stettner H, Beyer AS, Feigl G, Likar R., Br J Anaesth. 2011 May;106(5):732-7.
7. Blockade of intra-articular adrenergic receptors increases analgesic demands for pain relief after knee surgery.
Kager I, Mousa SA, Sieper J, Stein C, Pipam W, Likar R., Rheumatol Int. 2011 Oct;31(10):1299-306.
8. Intensity and treatment of symptoms in 3,030 palliative care patients: a cross-sectional survey of the EAPC Research Network.
Laugsand EA, Kaasa S, de Conno F, Hanks G, Klestad P; Research Steering Committee of the EAPC. J Opioid Manag. 2009 Jan-Feb;5(1):11-21.

2011

1. From the nursing home to hospital and back again... A mixed methods study on hospital transfers from nursing homes.
Kada O, Brunner E, Likar R, Pinter G, Leutgeb I, Francisci N, Pfeiffer B, Janig H. Z Evid Fortbild Qual Gesundhwes. 2011;105(10):714-22.
2. Quality of postoperative pain therapy in Austria: national survey of all departments of anaesthesiology.
Kinstner C, Likar R, Sandner-Kiesling A, Hutschala D, Pipam W, Gustorff B. Anaesthetist. 2011 Sep;60(9):827-34. doi: 10.1007/s00101-011-1911-2.
3. Tonsillotomy and adenotonsillectomy in childhood. Study on postoperative pain therapy.
Platzer M, Likar R, Stettner H, Jost R, Wutti C, Leipold H, Breschan C. Anaesthetist. 2011 Jul;60(7):625-32.
4. Recommendations for the management of chronic pain by intrathecal ziconotide.
Tronnier V, Hofacker W, Ilias W, Kleinmann B, Koulousakis A, Kress HG, Likar R, Lux EA, Müller-Schwefe G, Schütze G, Uberall M. MMW Fortschr Med. 2010 Oct 14;152 Suppl 3:106-9.
5. Clonidine for remifentanil-induced hyperalgesia: a double-blind randomized, placebo-controlled study of clonidine under intra-operative use of remifentanil in elective surgery of the shoulder.
Schlimp CJ, Pipam W, Wolrab C, Ohner C, Kager HI, Likar R. Schmerz. 2011 Jun;25(3):290-5.
6. Consecutive, prospective case series of a new method for ultrasound-guided supraclavicular approach to the brachiocephalic vein in children.
Breschan C, Platzer M, Jost R, Stettner H, Beyer AS, Feigl G, Likar R. Br J Anaesth. 2011 May;106(5):732-7.
7. Topography of the lumbar sympathetic trunk in normal lumbar spines and spines with spondylophytes.
Feigl GC, Kastner M, Ulz H, Breschan C, Dreu M, Likar R. Br J Anaesth. 2011 Feb;106(2):260-5.
8. Analgesic efficacy of TENS therapy in patients with gonarthrosis. A prospective, randomised, placebo-controlled, double-blind study.
Gschiel B, Kager H, Pipam W, Weichert K, Likar R. Schmerz. 2010 Sep;24(5):494-500.
9. Current knowledge of buprenorphine and its unique pharmacological profile
Pergolizzi J, Aloisi AM, Dahan A, Filitz J, Langford R, Likar R, Mercadante S, Morlion B, Raffa RB, Sabatowski R, Sacerdote P, Torres LM, Weinbroum AA. Pain Pract. 2010 Sep-Oct;10(5):428-50.

10. Blockade of intra-articular adrenergic receptors increases analgesic demands for pain relief after knee surgery.
Kager I, Mousa SA, Sieper J, Stein C, Pipam W, Likar R.
Rheumatol Int. 2011 Oct;31(10):1299-306.
11. Subcutaneous target stimulation (STS) in chronic noncancer pain: a nationwide retrospective study.
Sator-Katzenschlager S, Fiala K, Kress HG, Kofler A, Neuhold J, Kloimstein H, Ilias W, Mozes-Balla EM, Pinter M, Loining N, Fuchs W, Heinze G, Likar R.
Pain Pract. 2010 Jul-Aug;10(4):279-86. Epub 2010 Mar 3.
12. Efficacy of glycopyrronium bromide and scopolamine hydrobromide in patients with death rattle: a randomized controlled study.
Likar R, Rupacher E, Kager H, Molnar M, Pipam W, Sittl R.
Wien Klin Wochenschr. 2008;120(21-22):679-83.
13. Neuropathic pain among hospital in-patients. Analysis of Austrian hospital discharge data.
Dorner T, Gustorff B, Likar R, Lawrence K, Schwarz F, Rieder A.
Schmerz. 2009 Feb;23(1):59-64.
14. Transdermal buprenorphine in the management of persistent pain - safety aspects.
Likar R.
Ther Clin Risk Manag. 2006 Mar;2(1):115-25.
15. Improvement of pain therapy in Austria.
Likar R, Bernatzky G.
Schmerz. 2007 Jun;21(3):277.
16. Pain experience and pain therapy of tumor patients in the view of general practitioners.
Janig H, Pipam W, Lastin S, Sittl R, Bernatzky G, Likar R.
Schmerz. 2005 Apr;19(2):97-108.
17. Management of postoperative pain in children.
Sittl R, Griessinger N, Koppert W, Likar R.
Schmerz. 2000 Oct;14(5):333-9.